



Urinary problems

Common urinary problems in women are urinary incontinence, overactive bladder (OAB), and urinary tract infections (UTIs).



SIGNS & SYMPTOMS

Urinary incontinence means you lose bladder control or can't store urine like you should. Although there are many types, the most common ones in women are stress incontinence and urge incontinence.

Stress Incontinence

Urine leaks out with a sudden rise in pressure in the abdomen. This can occur when you cough, sneeze, lift, jump, run or strain to pass stool.

Urge Incontinence

Urine is released before you can get to the toilet due to a sudden and intense urge to urinate.

Overactive Bladder

You urinate often (8 or more times during the day and at least 2 times during the night) and you have a sudden and urgent need to urinate.

Urinary Tract Infections

Bladder Infection Symptoms

- You urinate more often than usual. It burns or stings when you urinate.
- Your urine is bloody or cloudy.
- You have pain in the abdomen or over your bladder.
- Confusion or other change in mental status, especially if you are over age 70.

Kidney Infection Symptoms

- Fever and shaking chills.
- Nausea and vomiting.
- Pain in one or both sides of your mid back.

Sometimes, there are no symptoms with a UTI.

CAUSES & RISK FACTORS

Urinary Incontinence

Problems occur with bladder muscles and nerves that help you hold or release urine and structures that support the bladder. This can be due to many factors:

- Physical changes due to aging or injury.
- Pregnancy and childbirth.
- Menopause.
- Multiple sclerosis.
- Spinal cord injury.
- Being overweight.

Overactive Bladder

Abnormal nerves send signals to the bladder at the wrong time. This causes spasms in the bladder muscles to squeeze without warning.

Urinary Tract Infections

Bacteria infect any part of the urinary tract – the bladder, kidneys and ureters (tubes that connect the kidneys to the bladder).





TREATMENT

Incontinence

- Bladder training, pelvic floor muscle training, or Kegel exercises.
- Medications.
- Medical treatment, such as an electric or magnetic stimulation device.
- Surgical procedures.

Overactive Bladder

Medications that help relax muscles of the bladder and prevent bladder spasms.

Urinary Tract Infections

An antibiotic is prescribed to treat the specific infection. Pain relievers are taken as needed.



TRIAGE QUESTIONS



Does a loss of bladder control come with any of these problems?

- A spine or back injury.
- Slurred speech.
- Loss of sight, double or blurred vision.
- Sudden, severe headache.
- Confusion and dizziness.
- Paralysis, weakness, or loss of feeling in an arm, leg, and/or the face on the same side of the body.

NO **YES** ➔ **GET EMERGENCY MEDICAL CARE**



Do you have kidney infection symptoms?

NO **YES** ➔ **GET EMERGENCY MEDICAL CARE**

CONTINUE IN NEXT COLUMN



Did you lose bladder control after a recent abdominal surgery or injury?

NO **YES** ➔ **SEE DOCTOR**



Do you have bladder infection symptoms?

NO **YES** ➔ **SEE DOCTOR**



Do you leak urine often or have constant dribbling of urine (signs of overflow incontinence)?

NO **YES** ➔ **SEE DOCTOR**



Do you have any of these problems?

- Constant urge to urinate.
- It feels like your bladder is still full after you urinate.
- Your urine smells bad.
- It hurts to have sex.

NO **YES** ➔ **SEE DOCTOR**



Have you had more than 3 bladder infections within 6 months or more than 4 bladder infections in the same year?

NO **YES** ➔ **SEE DOCTOR**



Did you get a skin rash or vaginal yeast infection after taking medicine prescribed for a urinary tract infection? Or, have symptoms not cleared up after taking prescribed medicine for 3 days?

NO **YES** ➔ **CALL DOCTOR**



Do you leak urine after you jump, cough, sneeze, laugh or lift heavy objects? Do you want to find out about treatment(s) for loss of bladder control?

NO **YES** ➔ **CALL DOCTOR**



USE SELF-CARE



SELF-CARE

Urinary Tract Infections

- Drink at least 8 glasses of water a day. Unsweetened cranberry juice may also help.
- For pain, take ibuprofen, acetaminophen, naproxen sodium, or Uristat®, an over-the-counter medicine for bladder infection pain.
- Wear cotton underwear and loose-fitting slacks. Use a heating pad on your abdomen.
- Avoid alcohol, caffeine and spicy foods.



Urinary Incontinence

- Avoid caffeine. Limit carbonated drinks, alcohol, citrus juices, greasy and spicy foods, and artificial sweeteners. Limit or avoid fluids 2 to 3 hours before bedtime.
- Empty your bladder before you leave the house, take a nap, or go to bed. Wear absorbent pads or briefs, as needed.
- Track how often you go to the bathroom, then slowly add 15 minutes between bathroom visits. Gradually increase the amount of time between visits to train your bladder to hold more urine before you need to go again.
- Keep a diary of when you leak urine. If you do this every 3 hours, empty your bladder every 2 hours. Use an alarm to remind you.
- Ask your doctor if your type of incontinence could be managed by a self-catheter or vaginal pessary. This small device supports pelvic floor muscles. A doctor needs to prescribe these devices.

Kegel Exercises are Pelvic Floor Exercises

These help treat or cure stress incontinence. Follow these steps:

1. Start to urinate, then hold back and try to stop. If you can slow the stream of urine, even a little, you are using the right muscles. You should feel muscles squeezing around the anus and the urethra (the tube through which urine is passed).
2. Relax your body. Close your eyes. Imagine that you are going to pass urine and then hold back from doing so. You should feel the muscles squeeze like you did in step 1.
3. Squeeze the muscles for 3 seconds. Then relax them for 3 seconds. When you squeeze and relax, count slowly. Start out doing this 3 times a day. Gradually work up to 3 sets of 10 contractions. Hold each one for 10 seconds at a time. You can do Kegel exercises when you lie down, sit and/or stand.
4. When you do these exercises do not tense the muscles in your belly or buttocks, hold your breath, clench your fists or teeth, or make a face.
5. Squeeze your pelvic floor muscles right before and during whatever it is (jumping, etc.) that causes you to leak urine. Relax the muscles once the activity is over.
6. You can also use pelvic weights prescribed by your doctor. You insert a weighted cone into the vagina and squeeze the correct muscles to keep it from falling out.

Do pelvic floor muscles daily. It may take several months to benefit from them.

FYI: INTERSTITIAL CYSTITIS (IC)

Note: Symptoms of a condition called Interstitial Cystitis (IC) mimic those of an acute UTI. Intense pain and pressure in the lower abdomen come with the need to urinate. (This can be more than 50 times a day.) Nine out of 10 persons who have IC are women. Antibiotics do not give relief, because bacteria is not present with IC. This condition needs medical diagnosis and treatment.



FOR MORE INFORMATION:

National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK)
niddk.nih.gov

National Association for Continence (NAFC)
800-BLADDER (252-3337)
nafc.org